



1

2

3 **biblio.ugent.be**

4

5 The UGent Institutional Repository is the electronic archiving and dissemination platform for all
6 UGent research publications. Ghent University has implemented a mandate stipulating that all
7 academic publications of UGent researchers should be deposited and archived in this repository.
8 Except for items where current copyright restrictions apply, these papers are available in Open
9 Access.

10

11 This item is the archived peer-reviewed author-version of:

12 Effect of temperature on the functional response of *Adalia bipunctata* to *Myzus persicae*

13 Jalali, M. A., L. Tirry & P. De Clercq

14 In: *BioControl*, 55: 261-269, 2010

15

16 **To refer to or to cite this work, please use the citation to the published version:**

17 Jalali, M. A., L. Tirry & P. De Clercq (2010). Effect of temperature on the functional response of
18 *Adalia bipunctata* to *Myzus persicae*. *BioControl* 55: 261-269. [10.1007/s10526-009-9237-6](https://doi.org/10.1007/s10526-009-9237-6)

19

Effect of temperature on the functional response of *Adalia bipunctata* to *Myzus persicae*

Jalali, M. A., L. Tirry & P. De Clercq

Department of Crop Protection, Ghent University, Coupure Links 653, B-9000 Ghent, Belgium ;

Patrick.DeClercq@ugent.be

Abstract

The effect of temperature on the functional response of female adults of the two-spot ladybird, *Adalia bipunctata* L. (Coleoptera: Coccinellidae) was examined in petri dish arenas containing sweet pepper leaves infested with different densities of the green peach aphid, *Myzus persicae* (Sulzer) (Hemiptera: Aphididae). The predator showed a type II functional response at three tested temperatures ranging from 19 to 27°C. The theoretical maximum number of prey captured by the predator increased with temperature. Based on the random predator equation, the estimated attack rates ranged from 0.13 h⁻¹ at 19°C to 0.35 h⁻¹ at 27°C on a leaf area of 20-25 cm². There was no significant difference between the attack rates of the predator at 23 and 27°C. Handling time significantly decreased as temperature increased from 19 (0.39 h) to 27°C (0.24 h). This study shows that *A. bipunctata* displays high predation rates on *M. persicae* for a wide range of temperatures, indicating its potential for augmentative releases against this aphid pest. The limitations of the predictions generated by functional response experiments are discussed.

41 **Keywords** *Adalia bipunctata*, *Myzus persicae*, Coccinellidae, Aphididae, functional response,
42 predation, augmentative biological control

43

44 **Running head:** Functional response of *Adalia bipunctata* on *Myzus persicae*

45

Introduction

A quantitative and qualitative understanding of the interaction among different species is crucial for the management of agricultural pests. Each species in an agricultural ecosystem is part of an often complex community, where it has to interact with many other species. These interactions may be competitive, predatory, parasitic or mutualistic (Begon et al. 1996; Bellows and Hassell 1999). One of the most important interactions with implications for biological control is the predator-prey relationship. How predators respond to changes in prey availability (i.e. functional response) is an issue of particular interest (Jeschke et al. 2002) and can determine whether a predator is able to regulate the density of its prey (Murdoch and Oaten 1975). The functional response has traditionally been defined as the number of prey attacked by an individual natural enemy in relation to host density over a given time interval (Solomon 1949). The response is characterized by an increase in the number of prey (or hosts) attacked over the lower range of prey densities, and an upper limit to the number of prey attacked at higher prey densities due to satiation in the case of predators (Mills 1982; Jeschke et al. 2002) and egg limitation or handling time constraints in the case of parasitoids (Getz & Mills 1996; Mills and Lacan 2004). Basically, functional responses have been delimited to three different types (Holling 1959). A type I response is characterized by a linear increase, a type II response by a monotonic decelerating increase, and a type III response by a sigmoidal increase in host numbers attacked. The type II functional response is the most frequently reported type in insects (Hassell et al. 1976; Begon et al. 1996), including coccinellids (Hodek and Honek 1996; Dixon 2000). However, predators may show different types of functional response curves as a function of host plant (Messina and Hanks 1998; De Clercq et al. 2000) or prey type (Sarmiento et al. 2007). *Harmonia axyridis*

(Pallas) has been reported to exhibit type I (Lou 1987), type II (He et al. 1994), and type III (Hu et al. 1989) functional responses on different aphid species. Interestingly, third instars and adults of *H. axyridis* exhibited different types of responses (type II and I, respectively) on the eggs of the monarch butterfly, *Danaus plexippus* (L.) (Koch et al. 2003).

In order to model predation by a long-lived, polyphagous predator, it is essential to know how the key parameters of the functional response (i.e. the attack coefficient, a , and handling time, T_h) scale with temperature (Thompson 1978). However, there are few data showing the influence of temperature on type and magnitude of the functional response of coccinellid predators. Overall, the predation rate of larval and adult stages of coccinellids has been reported to increase with temperature within a range of ca. 10-30 °C (Mack and Smilowitz 1982a,b; Giroux et al. 1995; Xia et al. 2003; Gotoh et al. 2004).

The two-spotted ladybird beetle, *Adalia bipunctata* (L.), is a common natural enemy of aphids in Europe and Central Asia (Mills 1979; Majerus 1994) and has been considered a potential biological control agent in different agro-ecosystems (Majerus 1994; Hodek and Honek 1996; Dixon 2000; Omkar and Pervez 2005). Although some studies have investigated the functional response of *A. bipunctata* (Mills 1979, 1982; Hemptinne et al. 1996), none of them have addressed the effect of temperature on the functional response of the predator. Mills (1979, 1982) found that prey consumption by female adults of *A. bipunctata* increased curvilinearly suggesting a type II functional response. On the contrary, the males of this species were not found to show a functional response (Hemptinne et al. 1996).

The objective of the current study is to investigate the effects of temperature on the functional response of *A. bipunctata* to different densities of the green peach aphid, *Myzus persicae* (Sulzer). Improving our understanding of the predator-prey interactions between *A.*

bipunctata and *M. persicae* may be helpful to optimize biological control of this economically important aphid pest.

Materials and Methods

Insects

Adults of *A. bipunctata* were taken from a colony at the Laboratory of Agrozoology, Ghent University, which was started with larvae purchased from Biobest NV (Westerlo, Belgium). At the commercial facility, the ladybird had been fed with live pea aphids, *Acyrtosiphon pisum* (Harris). Upon introduction at the Laboratory of Agrozoology, the predator was reared for one generation on an ad libitum supply of different nymphal stages of the green peach aphid, *M. persicae*, at each of the examined temperatures before starting the functional response study. The aphids were reared on sweet pepper plants (*Capsicum annuum* L., cv. California Wonder) in a culture room at $25 \pm 2^\circ\text{C}$, $50 \pm 10\%$ RH, and a photoperiod of 16:8 (L:D) h.

Functional response experiment

The experiment was performed at three constant temperatures (19, 23 and $27 \pm 1^\circ\text{C}$), a relative humidity of $65 \pm 5\%$ and a photoperiod of 16:8 (L:D) h. The selected temperatures should reflect thermal conditions frequently experienced by the predator in different protected and field crops in temperate areas. The female *A. bipunctata* used in the experiment had body weights ranging from 12.1-13.4 mg. At each of the tested temperatures, females were used that had initiated oviposition (i.e. 8-10 days old). To standardize the response, predators were starved for 24 h in boxes containing only a source of moisture before release into the test arena. The experimental arena consisted of a plastic Petri dish (14 cm in diameter) lined with paper towel. Each dish contained a sweet pepper leaf with its petioles inserted into an Eppendorf tube (2ml) filled with water. The average leaf area (both sides) was estimated to be approximately 20-25 cm² (n = 5 leaves). Third and fourth instars of *M. persicae* were used as prey in the experiment. Aphids were gently transferred by a fine camel hair brush from plants of the stock culture to the leaves in the test arenas. Prey densities used were: 5, 10, 15, 20, 30, 60, 80, 100 and 130 aphids per leaf. The aphids were allowed to settle and a female adult predator was introduced in each Petri dish. At each prey density, there were ten replicates for predator treatments and 5 controls (i.e. arenas without a predator). The total number of prey killed during a 24-h period was recorded. Killed prey were not replaced during the experiment (prey depletion method). Predation mortality was corrected for control mortality using the formula given by Xia et al. (2003):

$$Ne = N_0 \frac{N_d - N_c}{N_0 - N_c} \quad (1),$$

where N_e represents the number of prey killed, N_0 represents the initial number of prey, N_d represents the number of prey eaten and dead in the treatment, and N_c represents the number of prey dead in the control.

Data analysis

Data were analysed using SPSS (SPSS Inc. 2006) for analysis of variance (ANOVA) and SAS (SAS Institute 1999) for analysis of functional response. Data were submitted to a two-way ANOVA for the significance of the main effects of prey density and temperature on predation and their interaction.

Analysis of the functional response consists of two distinctive steps (De Clercq et al. 2000; Juliano 2001): determination of the type of functional response and estimation of the parameters of the functional response curve. The shape of the functional response is determined by a logistic regression of the proportion of prey eaten as a function of initial density. This is done by fitting a polynomial function:

$$\frac{N_e}{N_0} = \frac{\exp(P_0 + P_1 N_0 + P_2 N_0^2 + P_3 N_0^3)}{1 + \exp(P_0 + P_1 N_0 + P_2 N_0^2 + P_3 N_0^3)} \quad (2),$$

where N_e is the number of prey eaten, N_0 is the initial number of prey, and P_0 , P_1 , P_2 , and P_3 are the parameters to be estimated. To estimate these parameters, the CATMOD procedure was used (Juliano 2001). The data sets were individually fitted to the model (2), for determining the type of functional response (type II or III). The sign of P_1 and P_2 can be used to distinguish the shape of the curves. A positive linear parameter (P_1) and a negative quadratic parameter (P_2) indicate that the functional response is type III, whereas if both parameters are negative, the functional response is type II (Juliano 2001). In the second step, a nonlinear least square regression (NLIN) procedure was used to estimate the parameters of the random predator

equation (T_h and a for type II and b , c , and d for type III) (Rogers 1972; Juliano 2001). These equations are described by:

$$N_e = N_0 \{ 1 - \exp[a(T_h N_e - T)] \} \quad (\text{type II, prey depletion}) \quad (3),$$

$$N_e = N_0 \{ 1 - \exp[(d + bN_0)(T_h N_e - T)/(1 + cN_0)] \} \quad (\text{type III, prey depletion}) \quad (4),$$

where N_e is the number of prey eaten, N_0 the initial number of prey, a the attack constant, T the total time available (24 h) and T_h the handling time per prey, and b , c , and d are constants. Lower densities where prey were completely depleted by the end of the experiment were omitted from the analysis to avoid biased estimates (as discussed below). Separation of statistically different parameter estimates was done using asymptotic 95% confidence intervals (CI) (Juliano 2001).

Results

The proportion of prey consumed by the predator declined with increasing prey density, justifying the decision to use non-linear curve fitting (Table 1). The negative linear coefficient ($P_1 < 0$) suggests that there was a type II functional response at all tested temperatures. Because estimates of the linear coefficient in the original cubic model were not significantly different from 0 ($P > 0.05$) in the case of 23°C and 27°C, the model was reduced by subsequently omitting the cubic term until all remaining coefficients were significant (Juliano, 2001; Mahdian et al., 2006). The negative sign of the estimates of the linear coefficient (P_1) in the reduced models for

23°C (quadratic model: $P_1 = -0.0884$, $P < 0.0001$) and 27°C (quadratic model: $P_1 = -0.0869$, $P < 0.0001$) confirmed the designated types of functional response in the original model. Significant quadratic terms (P_2) were derived for *A. bipunctata* at temperatures ranging from 19°C and 23°C, indicating that asymptotically declining proportions of prey were killed by the predator at these temperatures.

All data were fitted to the random predator equation. The coefficients of determination (R^2) indicated greater variation in predation rates at 23 °C than at 19 and 27°C (Table 2). The number of *M. persicae* nymphs killed by *A. bipunctata* females increased at a decreasing rate until reaching an upper plateau, as the number of *M. persicae* offered increased (Fig. 1). The theoretical maximum number of prey captured by the predator increased with temperature. A two-way ANOVA with prey density and temperature as factors revealed a significant difference in predation at the different prey densities ($F_{8,243} = 878.90$, $P < 0.001$) and across temperatures ($F_{2,243} = 185.54$, $P < 0.001$). The interaction between the two factors was also significant ($F_{16,243} = 30.54$, $P < 0.001$).

The relationship between temperature, handling times and attack rates, calculated from the response curves, is shown in Table 2. Handling time significantly decreased ($F_{2,29} = 344.083$, $P < 0.001$) as temperature increased from 19 (0.39h) to 27°C (0.24h). Likewise, there was a significant effect of temperature on attack rate ($F_{2,29} = 26.414$, $P < 0.001$). Attack rate on a 20-25 cm² leaf area at 23°C (0.31 hour⁻¹) was higher than that at 19°C (0.13 hour⁻¹), but similar to that at 27°C (0.35 hour⁻¹). The estimated maximum numbers of 3rd-4th instars of *M. persicae* that can be killed by a single *A. bipunctata* female (T/T_h) were 2.56, 2.78 and 4.17 per hour, at 19, 23 and 27°C, respectively.

198

199 **Discussion**

200

201 Juliano (2001) and Tully et al. (2005) reviewed experimental circumstances that violate the basic
202 assumptions of Holling's (1959) disc equation, which the most frequently used equation to
203 model functional responses of predatory arthropods. One of the most common violations is prey
204 depletion during the experiment. When killed prey are not replaced during exposure to the
205 predator, Rogers' (1972) model can be used as it accounts for changing prey densities. However,
206 this model cannot correct for complete prey depletion before the end of the experiment. When
207 all prey are eaten within the observation interval, no valid information can be derived from the
208 experiment and including these densities in the analysis may yield biased estimates. Therefore,
209 lower prey densities where prey were completely depleted by the end of the experiment were
210 removed from the functional response analysis in the present study. Ideally, either the arena
211 size should be increased or prey should be replenished at shorter time intervals in order to avoid
212 complete prey depletion at lower initial densities of prey.

213 Between 19 and 27°C, the rate of predation by adult females of *A. bipunctata* on 3rd and
214 4th instars of *M. persicae* increased with increasing prey density, but the mean ratio of prey
215 killed decreased monotonically, suggesting a type II functional response (Holling 1959, 1965;
216 Gotelli 1995). A similar response curve was reported earlier by Mills (1982) for *A. bipunctata*
217 preying on *A. pisum* at 20°C. Type II functional response curves have been reported in a number
218 of other coccinellids (Uygun and Atlihan 2000; Omkar and Pervez 2003; Lee and Kang 2004;
219 Pervez and Omkar 2005; Mandour et al. 2006; Moura et al. 2006). However, type III functional

responses have been also reported for coccinellids (Hu et al. 1989; Messina and Hanks 1998; Sarmento et al. 2007).

Some authors have challenged the efficacy of predators showing type II functional responses as biological control agents. Holling (1965) stated that predators showing a type III response are theoretically more capable of suppressing prey populations. Although the shape of the functional response curve is an important factor, it is insufficient as a criterion to predict success or failure of a predator as a biocontrol agent, as other factors such as numerical response, intrinsic growth rates, host patchiness, competition, and environmental complexities (abiotic and biotic factors) also have a major influence on the efficiency of a predator in managing a pest population (Pervez and Omkar 2005; see also below). A review by Fernández-arhex and Corley (2003) showed that most parasitoid species have a type II response, but there is no clear relationship between curve shape and biocontrol success in these insects. The form of the functional response curve may vary according to environmental conditions (Abrams 1982), such as habitat heterogeneity (Hildrew and Townsend 1977; Kaiser 1983; Lipcius and Hines 1986), light intensity (Koski & Johnson 2002), host plant (Carter et al. 1984; Messina and Hanks 1998; De Clercq et al. 2000; Skirvin and Fenlon 2001), temperature (Thompson 1978; Gresens et al. 1982; Bailey 1989; Mohaghegh et al. 2001; Mahdian et al. 2006; Li et al. 2007) and refuge for prey (Lipcius and Hines 1986; Messina and Hanks 1998). In the current study, temperature clearly influenced the magnitude of predator response to prey density (Fig.1). This may be attributed to temperature-related changes in the metabolism and activity of predator and prey (McCoull et al. 1998). Higher temperatures increased the functional response of *A. bipunctata* by increasing attack rate and decreasing handling time. Similar observations were reported by Thompson (1978), Gresens et al. (1982), Bailey (1989), Anderson et al. (2001) and Gotoh et al. (2004). Our findings suggest that *A. bipunctata* spent more time in non-searching activities (e.g.

resting) at low temperatures (19°C), while more intensive searching and feeding occurred at higher temperatures (above 23°C).

The current study suggests that an individual *A. bipunctata* female can kill a theoretical maximum number of 61, 67 and 100 3rd-4th instars of *M. persicae* per day at 19, 23 and 27°C, respectively. These numbers are higher than those reported by Chi and Yang (2003) for *Propylea japonica* Thunberg and Cabral et al. (2006) for *Coccinella undecimpunctata* L. preying on *M. persicae*. However, comparison of the predation rates of *A. bipunctata* in our study with those of other coccinellids is complicated by variability in the size of predators and prey and in experimental conditions (e.g., test arena, temperature).

Functional responses may provide important information on the voracity of a biological control agent, and on the effects of abiotic (e.g., temperature) or biotic (e.g., host plants) factors on its foraging efficiency (e.g., Carter et al. 1984; Messina and Hanks 1998; De Clercq et al. 2000; Mohaghegh et al. 2001; Skirvin and Fenlon 2001; Mahdian et al. 2006; Li et al. 2007). However, functional response studies have been criticized because they are often performed in small artificial arenas using unrealistic prey densities and do not consider spatial habitat complexities or multispecies prey situations (van Lenteren and Bakker 1976; Murdoch 1983; O'Neil 1989, 1997; Kareiva 1990; Wiedenmann and O'Neil 1991; Messina and Hanks 1998; Hardman et al. 1999). Furthermore, functional responses do not consider crucial life history parameters of a predator that may affect its value as a biological control agent. In the case of coccinellid predators, the selection of prey patches by female adults is determined by expectations of future bottlenecks in prey abundance (Dixon 2000). Many aphidophagous predators have evolved the ability to attack prey colonies at an early stage of development to maximize the fitness of their offspring. Dixon (2000) proposed that the most important attribute of an

effective biological control agent is the rate of its development relative to that of its prey, i.e. the generation time ratio. According to this hypothesis, if the developmental rate of a predator is similar to or faster than that of the prey then the predator is potentially capable of suppressing the prey population. This concept would render *A. bipunctata* and other aphidophagous coccinellids ineffective biological control agents, given the high intrinsic rates of increase of most aphid species (Dixon 2000). However, advanced larval stages are released in relatively high numbers in many augmentative biological control programmes using aphidophagous coccinellids (e.g., Ferran et al. 1998; Wyss et al. 1999a,b). Commercial packagings of *A. bipunctata* usually contain second to third instars (J. Vermeulen, BioBest NV, personal communication). In such an inundative approach the objective is short term pest suppression by the released larvae and the resulting adults rather than long term control of the aphid populations. In addition, *A. bipunctata* larvae can be released in aphid hot spots rather than in the whole crop.

The current study shows that *A. bipunctata* displays relatively high predation rates on *M. persicae* at a wide range of temperatures, indicating its potential for augmentative releases against the green peach aphid. Functional response models have served as a basis for developing ecological theory on foraging behaviour and tritrophic interactions between plants, prey and predators (or parasitoids) (Holling 1959; Hassell 1978; Messina and Hanks 1998; Tully et al. 2005). Measures of voracity from functional response experiments may be indicative of the value of coccinellid predators as augmentative biological control agents and provide a basis for determining release rates appropriate for various aphid densities. However, robust experimental assessment of functional responses remains difficult and certain behavioural responses that are crucial in a biological control context cannot be taken into account in the experimental design (van Lenteren and Bakker 1976, 1978; Tully et al. 2005). For instance,

291 laboratory studies may indicate the predation potential of *A. bipunctata* when aphids are
292 abundant, but they do not consider cannibalism or emigration of the predator that may occur
293 when aphid populations start to decline, leaving few predators in the crop in case the aphids
294 would resurge. Therefore, more field studies are needed to understand the foraging behaviour
295 of *A. bipunctata* in different cropping systems, in order to design practical release strategies for
296 this coccinellid.

297

298 **Acknowledgments**

299

300 We are grateful to Steven A. Juliano, Isabelle Vantornhout, Kamran Mahdian and two
301 anonymous reviewers for their helpful advice. M. A. Jalali was supported by PhD grant no.
302 810086 from the Ministry of Science, Research and Technology of Iran.

303

304

305

References

- Abrams P (1982) Functional responses of optimal foragers. *Am Nat* 120: 382–390
- Anderson MT, Kiesecker JM, Chivers DP, Blaustein AR (2001) The direct and indirect effects of temperature on a predator–prey relationship. *Can J Zool* 79: 1834–1841
- Bailey PCE (1989) The effect of water temperature on the functional response of the water stick *Ranatra dispar* (Heteroptera: Nepidae). *Aust J Ecol* 14: 381–386
- Begon M, Harper JL, Townsend CR (1996) Ecology: individuals, populations, and communities. Blackwell, London
- Bellows TS, Hassell MP (1999) Theories and mechanisms of natural population regulation. In: Bellows TS, Fisher TW (eds) Handbook of biological control. Academic Press, San Diego, pp 17–44
- Cabral S, Soares AO, Moura R, Garcia P (2006) Suitability of *Aphis fabae*, *Myzus persicae* (Homoptera: Aphididae) and *Aleyrodes proletella* (Homoptera: Aleyrodidae) as prey for *Coccinella undecimpunctata* (Coleoptera: Coccinellidae). *Biol Control* 39: 434–440
- Carter MC, Sutherland D, Dixon AFG (1984) Plant structure and the searching efficiency of coccinellid larvae. *Oecologia* 63: 394–397
- Chi H, Yang TC (2003) Two-sex life table and predation rate of *Propylaea japonica* Thunberg (Coleoptera: Coccinellidae) fed on *Myzus persicae* (Sulzer) (Homoptera: Aphididae). *Environ Entomol* 32: 327–333

- 326 De Clercq P, Mohaghegh J, Tirry L (2000) Effect of host plant on the functional response of the
327 predator *Podisus nigrispinus* (Heteroptera: Pentatomidae). Biol Control 18: 65–70
- 328 Dixon AFG (2000) Insect predator–prey dynamics. Ladybird beetles & biological control.
329 Cambridge University Press, Cambridge.
- 330 Ferran A, Giuge L, Tourniaire R, Gambier J, Fournier D (1998) An artificial non-flying mutation to
331 improve the efficiency of the ladybird *Harmonia axyridis* in biological control of aphids.
332 BioControl 43: 53-64
- 333 Fernández-arhex V, Corley JC (2003) The functional response of parasitoids and its implications
334 for biological control. Biocontrol Sci Technol 13: 403-413
- 335 Getz WM, Mills NJ (1996) Host–parasitoid coexistence and egg-limited encounter rates. Am Nat
336 148: 301–315
- 337 Giroux S, Duchesne RM, Coderre D (1995) Predation of *Leptinotarsa decemlineata* (Coleoptera,
338 Chrysomelidae) by *Coleomegilla maculata* (Coleoptera, Coccinellidae) – Comparative
339 effectiveness of predator developmental stages and effect of temperature Environ
340 Entomol 24: 748-754
- 341 Gotelli NJ (1995) A primer of ecology. Sinauer Associates, Inc., Sunderland, Massachusetts.
- 342 Gotoh T, Nozawa M, Yamauchi K (2004) Prey consumption and functional response of three
343 acarophagous species to eggs of the two-spotted spider mite in the laboratory. Appl
344 Entomol Zool 39: 97–105

- 345 Gresens SE, Cothran ML, Thorp JH (1982) The influence of temperature on the functional
346 response of the dragonfly *Celithemis fasciata* (Odonata: Libellulidae). *Oecologia* 53: 281–
347 284
- 348 Hardman JM, van der Werf W, Nyrop JP (1999) Modeling mite dynamics on apple trees in
349 eastern North America. *Proceedings 5th International Symposium on Computer*
350 Modelling in Fruit Research and Orchard Management. *Acta Horti* 499: 201-209
- 351 Hassell MP (1978) The dynamics of arthropod predator–prey systems. Princeton University
352 Press, Princeton
- 353 Hassell MP, Lawton JH, Beddington, JR (1976) Components of arthropod predation. 1. Prey
354 death rate. *J Anim Ecol* 45: 135–164
- 355 Hassell MP, Lawton JH, Beddington, JR (1977) Sigmoid functional response by invertebrate
356 predators and parasitoids. *J Anim Ecol* 46: 249–262
- 357 He JL, Ma EP, Shen YC, Chen WL, Sun XQ (1994) Observations of the biological characteristics of
358 *Harmonia axyridis* (Pallas) (Coleoptera: Coccinellidae). *J Shanghai Agric Coll* 12: 119–124
- 359 Hemptinne J-L, Dixon AFG, Lognay G (1996) Searching behaviour and mate recognition by males
360 of the two-spot ladybird beetle, *Adalia bipunctata*. *Ecol Entomol* 21: 165–170
- 361 Hildrew AG, Townsend CR (1977) The influence of substrate on the functional response of
362 *Plectrocnemia conspersa* (Curtis) larvae (Trichoptera: Polycentropodidae). *Oecologia* 31:
363 21–26
- 364 Hodek I, Honek A (1996) Ecology of Coccinellidae. Kluwer Academic Publishers, Dordrecht

365 Holling CS (1959) Some characteristics of simple types of predation and parasitism. Can Entomol
 366 91: 385–398

367 Holling CS (1965) The functional response of predators to prey density and its role in mimicry
 368 and population regulation. Mem Entomol Soc Can 48: 3–60

369 Hu YS, Wang ZM, Ning CL, Pi ZQ, Gao GQ (1989) The functional response of *Harmonia (Leis)*
 370 *axyridis* to their prey of *Cinara* sp. Nat Enemies Insects 11: 164–168

371 Jeschke JM, Kopp M, Tollrian R (2002) Predator functional responses: discriminating between
 372 handling and digesting prey. Ecol Monog 72: 95–112

373 Juliano SA (2001) Nonlinear curve fitting: predation and functional response curves. In: Cheiner
 374 SM, Gurven J (eds) Design and analysis of ecological experiments, 2nd edn. Chapman &
 375 Hall, New York, pp 159–182

376 Kaiser H (1983) Small scale spatial heterogeneity influences predation success in an unexpected
 377 way: model experiments on the functional response of predatory mites (Acarina).
 378 Oecologia 56: 249–256

379 Kareiva P (1990) Population dynamics in spatial complex environments: theory and data. Philos.
 380 Trans R Soc London B 330: 175–190

381 Koch RL, Hutchison WD, Venette RC, Heimpel GE (2003) Susceptibility of immature monarch
 382 butterfly, *Danaus plexippus* (Lepidoptera: Nymphalidae: Danainae), to predation by
 383 *Harmonia axyridis* (Coleoptera: Coccinellidae). Biol Control 28: 265–270

384 Koski ML, Johnson BM (2002) Functional response of kokanee salmon (*Oncorhynchus nerka*) to
 385 *Daphnia* at different light levels. Can J Fish Aquat Sci 59: 707–716

- 386 Lee JH, Kang TJ (2004) Functional response of *Harmonia axyridis* (Pallas) (Coleoptera:
387 Coccinellidae) to *Aphis gossypii* Glover (Homoptera: Aphididae) in the laboratory. Biol
388 Control 31: 306-310
- 389 Li DX, Tian J, Shen ZR (2007) Functional response of the predator *Scolothrips takahashii* to
390 hawthorn spider mite, *Tetranychus viennensis*: effect of age and temperature.
391 BioControl 52: 41-61
- 392 Lipcius RN, Hines AH (1986) Variable function responses of a marine predator in dissimilar
393 homogeneous microhabitats. Ecology 67: 1361–1371
- 394 Lou HH (1987) Functional response of *Harmonia axyridis* to the density of *Rhopalosiphum*
395 *prunifoliae*. Nat Enemies Insects 9: 84-87
- 396 Mack TP, Smilowitz Z (1982a) Using temperature-mediated functional response models to
397 predict the impact of *Coleomegilla maculata* (DeGeer) adults and third instar larvae on
398 green peach aphids. Environ Entomol 11: 46–52
- 399 Mack TP, Smilowitz Z (1982b) CMACSIM, a temperature dependent predator-prey model
400 simulating the impact of *Coleomegilla maculata* (DeGeer) on green peach aphids on
401 potato plants. Environ Entomol 11: 1193–1201
- 402 Mahdian K, Vantornhout I, Tirry L, De Clercq P (2006) Effects of temperature on predation by the
403 stinkbugs *Picromerus bidens* and *Podisus maculiventris* (Heteroptera: Pentatomidae) on
404 noctuid caterpillars. Bull Entomol Res 96: 489–496
- 405 Majerus MEN (1994) Ladybirds. Harper Collins, London

406 Mandour NS, El-Basha NAS, Liu TX (2006) Functional response of the ladybird, *Cydonia vicina*
 407 *nilotica* to cowpea aphid, *Aphis craccivora* in the laboratory. Insect Sci 13: 49-54

408 McCoull CJ, Swain R, Barnes RW (1998) Effect of temperature on the functional response and
 409 components of attack rate in *Naucoris congrex* Stal (Hemiptera: Naucoridae). *Aust J*
 410 *Entomol* 37: 323–327

411 Messina FJ, Hanks JB (1998) Host plant alters the shape of the functional response of an aphid
 412 predator (Coleoptera: Coccinellidae). *Environ Entomol* 27: 1196–1202

413 Mills NJ (1979) *Adalia bipunctata* (L.) as a generalist predator of aphids. Ph.D. Thesis, University
 414 of East Anglia, Norwich, UK

415 Mills NJ (1982) Satiation and the functional response: a test of a new model. *Ecol. Entomol.* 7:
 416 305–315

417 Mills NJ, Lacan I (2004) Ratio dependence in the functional response of insect parasitoids:
 418 evidence from *Trichogramma minutum* foraging for eggs in small host patches. *Ecol*
 419 *Entomol* 29: 208–216

420 Mohaghegh J, De Clercq P, Tirry L (2001) Functional response of the predators *Podisus*
 421 *maculiventris* (Say) and *Podisus nigrispinus* (Dallas) (Het., Pentatomidae) to the beet
 422 armyworm, *Spodoptera exigua* (Hübner) (Lep., Noctuidae): effect of temperature. *J Appl*
 423 *Entomol* 125: 131-134

424 Moura R, Garcia P, Soares AO (2006) Does pirimicarb affect the voracity of the euriphagous
 425 predator, *Coccinella undecimpunctata* L. (Coleoptera: Coccinellidae)? *Biol Control* 38:
 426 363–368

- 427 Murdoch WW (1983) The functional response of predators. J Appl Ecol 10: 335–342
- 428 Murdoch WW, Oaten A (1975) Predation and population stability. Adv Ecol Res 9: 2–131
- 429 O’Neil RJ (1989) Comparison of laboratory and field measurements of the functional response of
430 *Podisus maculiventris* (Hemiptera: Pentatomidae). J Kans Entomol Soc 70: 40–48
- 431 O’Neil RJ (1997) Functional response and search strategy of *Podisus maculiventris* (Heteroptera:
432 Pentatomidae) attacking Colorado potato beetle (Coleoptera: Chrysomelidae). Environ
433 Entomol 26: 1183–1190
- 434 Omkar, Pervez A (2003) Influence of prey deprivation on biological attributes of pale morphs of
435 the ladybeetle, *Propylea dissecta* (Mulsant). Insect Sci Applic 23: 143–148
- 436 Omkar, Pervez A (2005) Ecology of two-spotted ladybird, *Adalia bipunctata*: a review. J Appl
437 Entomol 129: 465-474
- 438 Pervez A, Omkar (2005) Functional responses of coccinellid predators: An illustration of a logistic
439 approach. J Insect Sci 5: 1-6
- 440 Rogers D (1972) Random search and insect population models. J Anim Ecol 41: 369-383
- 441 Sarmento RA, Pallini A, Venzon M, DeSouza O, Molina-Rugama AJ, Oliveira CL (2007) Functional
442 response of the predator *Eriopis connexa* (Coleoptera: Coccinellidae) to different prey
443 types. Braz Arch Biol Technol 50: 121–126
- 444 SAS Institute (1999) SAS/STA User’s Guide, Version 1.3.0.161. SAS Institute, Cary, NC, USA

445 Skirvin DJ, Fenlon JS (2001) Plant species modifies the functional response of *Phytoseiulus*
 446 *persimilis* (Acari: Phytoseiidae) to *Tetranychus urticae* (Acari: Tetranychidae):
 447 implications for biological control. Bull Entomol Res 91: 61-67

448 Solomon ME (1949) The natural control of animal populations. J Anim Ecol 18: 1–35

449 SPSS Inc (2006) SPSS version 15.0. SPSS Inc, Chicago, IL

450 Thompson DJ (1978) Towards a realistic predator-prey model: The effect of temperature on the
 451 functional response and life history of larvae of the damselfly, *Ischnura elegans*. J Anim
 452 Ecol 47: 757-767

453 Tully T, Cassey P, Ferrière R (2005) Functional response: rigorous estimation and sensitivity to
 454 genetic variation in prey. Oikos 111: 479-487

455 Uygun N, Atlihan R (2000) The effect of temperature on development and fecundity of *Scymnus*
 456 *levailanthi*. BioControl 45: 453–462

457 van Lenteren JC, Bakker K (1976) Functional response in invertebrates. Neth J Zool 26: 567-572

458 van Lenteren JC, Bakker K (1978) Behavioural aspects of the functional responses of a parasite
 459 (*Pseudeucoila bochei* Weld) to its host (*Drosophila melanogaster*). Neth J Zool 28: 213-
 460 233

461 Wiedenmann RN, O'Neil RJ (1991) Laboratory measurement of the functional response of
 462 *Podisus maculiventris* (Say) (Heteroptera: Pentatomidae). Environ Entomol 20: 610–614

463 Wyss, E, Villiger M, Hemptinne J-L, Müller-Schärer H (1999a) Effects of augmentative releases of
 464 eggs and larvae of the two-spot ladybird beetle, *Adalia bipunctata*, on the abundance of

465 the rosy apple aphid, *Disaphis plantaginea*, in organic apple orchards. Entomol Exp Appl
466 90: 167-173

467 Wyss, E, Villiger M, Müller-Schärer H (1999b) The potential of three native insect predators to
468 control the rosy apple aphid, *Disaphis plantaginea*. BioControl 44: 171-182

469 Xia JY, Rabbinge R, van der Werf W (2003) Multistage functional responses in a ladybeetle-aphid
470 system scaling up from the laboratory to the field. Environ Entomol 32: 151–162

471

Table 1 Results of logistic regression analysis of the proportion of *M. persicae* nymphs (3rd and 4th instars) eaten by adult females of *A. bipunctata* on initial prey density at different temperatures

Temperature (°C)	Coefficient	Estimate	SE	Chi Square	P-value
19	Constant (P_0)	6.7094	0.8607	60.7704	<0.0001
	Linear (P_1)	-0.1911	0.0336	32.3628	<0.0001
	Quadratic (P_2)	0.0019	0.0004	20.1901	<0.0001
	Cubic (P_{C3})	-6.21E-6	1.591E-6	15.2199	<0.0001
23	Constant (P_0)	-3.7786	4.0637	0.8646	0.3524
	Linear (P_1)	0.2378	0.1375	2.9906	0.0837
	Quadratic (P_2)	-0.0032	0.0015	4.6605	0.0309
	Cubic (P_{C3})	0.12E-4	5.188E-6	5.6793	0.0172
27	Constant (P_0)	2.1165	6.3683	0.1105	0.7396
	Linear (P_1)	0.0910	0.2120	0.1844	0.6676
	Quadratic (P_2)	-0.0016	0.0023	0.5232	0.4695
	Cubic (P_{C3})	6.568E-6	7.798E-6	0.7096	0.3996

Table 2 Parameters estimated by the random predator equation and corresponding asymptotic 95% confidence intervals (CI) indicating functional response of adult females of *A. bipunctata* to densities of *M. persicae* 3rd-4th instars at different temperatures

Temperature (°C)	Type	R ²	a (h ⁻¹)*		T _h (h)	
			Mean ± SE	95% CI	Mean ± SE	95% CI
19	II	0.67	0.13±0.03	0.07-0.18	0.39±0.03	0.34-0.45
23	II	0.26	0.31±0.11	0.08-0.53	0.36±0.02	0.32-0.41
27	II	0.74	0.35±0.09	0.16-0.54	0.24±0.01	0.21-0.27

*a: attack rate (on a 20-25 cm² leaf area); T_h: handling time

485 **Fig. 1** Observed numbers of *M. persicae* 3rd-4th instars killed in 24h by *A. bipunctata* females vs.
486 numbers predicted by the random predator equation (lines) at three different temperatures on
487 a 20-25 cm² leaf area. Lines were only drawn through densities included in the analysis (i.e.
488 those without complete prey depletion).




Fig. 1.